## AUTHOR INDEX

ABBAS, S. A., 201
ABOLA, J., 1
ANGYAL, S. J., 151
ANISUZZAMAN, A. K. M., 258

BRIMACOMBE, J. S., 234

DAIS, P., 159
DAMM, J. B. L., 43
DUBEY, R., 201
DUTTON, G. G. S., 213

EL KHADEM, H. S., 13
ENNIFAR, S., 13

FAGHIH, R., 247
FRASER-REID, B., 247

GLAUDEMANS, C. P. J., 23

HASHIMOTO, H., 171
HORTON, D., 258

ISBELL, H. S., 13
ISHIDO, Y., 189

JEANLOZ, R. W., 221
JEFFREY, G. A., 1, 35

KABIR, A. K. M. S., 234
KAGEYAMA, S., 241
KAMERLING, J. P., 43
KOVÁČ, P., 23
KUMA-MINTAH, A., 213

LEHMANN, J., 53

MALUSZYNSKA, H., 35
MATHLOUTHI, M., 83
MATTA, K. L., 201

NAKABAYASHI, S., 221

ODIER, L., 151
OGAWA, T., 127
OK, K., 69
ONODA, T., 241

PAULSEN, H., 105
PERLIN, A. S., 159

SAITO, Y., 171
SAKAKIBARA, T., 189
SEUVRE, A.-M., 83
STEIGER, K.-M., 105
STEVENS, J. D., 151
SUDOH, R., 189

TACHIMORI, Y., (the late) 189
TAKAGI, Y., 69
TAKAHASHI, Y., 127
TAKAI, I., 189
THOMAS, R. L., 201
TSUCHIYA, T., 69, 241

UMEZAWA, H., 69, 241
UMEZAWA, S., 69, 241

VAN PELT, J., 43
VLIEGENTHART, J. F. G., 43

WARREN, C. D., 221
WATANABE, T., 171
WHISTLER, R. L., 252
WITCZAK, Z. J., 252

YAMAMOTO, A., 189
YEH, H. J. C., 23
YEON, Y., 1
YOSHIMURA, J., 171

ZISER, L., 53

## SUBJECT INDEX

2-Acetamido-2,6-dideoxy-L-galactose, synthesis of, 258
Adriamycinone, synthesis of antitumor-active 7-*O*-(2,6-dideoxy-2-fluoro-α-L-talopyranosyl)-daunomycinone and, 69
Aldehyde-group participation, the reaction of 2,3:6,7-di-*O*-isopropylidene-5-*O*-methanesulphonyl-L-*glycero*-D-*manno*-heptofuranose with sodium methoxide: an, 234
ω-Aminated 1-methoxyalkyl β-D-glucopyranosides as potential β-D-glucosidase inhibitors, syntheses of homologous, 53
1,6-Anhydro-β-D-glucopyranose (levoglucosan), a ready route to a functionalized anhydro derivative of, 247
1,6-Anhydrohexoses, selective deuteration over Raney nickel in deuterium oxide, 151

4,6-*O*-Benzylidene-2-*C*- and -3-*C*-*p*-tolylsulfonylhex-2-enopyranoside derivatives and a 2-*C*-*p*-tolylsulfonylhex-2-enitol derivative, preparation of, 189

Convenient synthesis of 6-deoxy-2-*C*-methyl-D-mannitol and 2-*C*-(hydroxymethyl)-D-mannitol, branched-chain alditols, 252
Crystal structure of the *C*-nucleoside, 1,3-dimethyl 8-β-D-ribofuranosyl xanthine monohydrate, 35
Crystal structures of octyl α-D-glucopyranoside monohydrate and hemihydrate: mesogenic structures with interdigitizing alkyl chains, 1
Cyclo{→6)-[α-D-Glc*p*-(1→4)]$_5$-α-D-Glc*p*-(1→}, total synthesis of cyclomaltooctaose and an isomer of cyclomaltohexaose, 127
Cyclomaltooctaose and an isomer of cyclomaltohexaose, cyclo{→6)-[α-D-Glc*p*-(1→4)]$_5$-α-D-Glc*p*-(1→}, total synthesis of, 127

5-*O*-(4-Deoxy-4-fluoro-β-D-mycaminosyl)tylonolide, synthesis of, 241
Deuteration over Raney nickel in deuterium oxide: 1,6-anhydrohexoses, selective, 151
7-*O*-(2,6-Dideoxy-2-fluoro-α-L-talopyranosyl)-daunomycinone and -adriamycinone, synthesis of antitumor-active, 69
1,3-Dimethyl 8-β-D-ribofuranosyl xanthine monohydrate, the crystal structure of the *C*-nucleoside, 35
Dolichyl ester of tetrasaccharide diphosphate, synthesis, 213
*Escherichia coli* capsular antigen K34, structure of, 189

D-Fructose in dimethyl sulfoxide, intramolecular hydrogen-bonding and solvation contributions to the relative stability of the β-furanose form of, 159
F.t.-i.r. spectra of oligo- and poly-nucleotides, 83
L-Fucosamine, *N*-acetyl-, synthesis of, 258
Functionalized anhydro derivative of 1,6-anhydro-β-D-glucopyranose (levoglucosan), a ready route to a, 247

L-Galactose, 2-acetamido-2,6-dideoxy-, synthesis of, 258
β-D-Gal*p*NAc-(1→4)-β-D-Gal*p*-(1→4)-D-GlcNAc, synthesis of, 105
β-D-Gal*p*-(1→4)-β-D-Glc*p*NAc-(1→3)-β-D-Gal*p*-(1→4)-D-Glc, synthesis of, 105
β-D-Gal*p*-(1→4)-β-D-Glc*p*NAc-(1→4)-β-D-Gal*p*-(1→4)-D-Glc, synthesis of, 105
*O*-β-D-Glc*p*NAc-(1→3)-*O*-β-D-Gal*p*-(1→4)-D-GlcNAc, synthesis of, 201
*O*-β-D-Glc*p*NAc-(1→3)-*O*-β-D-Gal*p*-(1→3)-α-D-Gal*p*NAcOC$_6$H$_4$NO$_2$-2, synthesis of, 201
*O*-β-D-Glc*p*NAc-(1→3)-*O*-β-D-Gal*p*-(1→3)-β-D-Gal*p*NAcOC$_6$H$_4$NO$_2$-4, synthesis of, 201
Glucopyranosides, synthesis and n.m.r. spectra of methyl 2-deoxy-2-fluoro- and 3-deoxy-3-fluoro-α- and β-D-, 23
β-D-Glucosidase inhibitors, syntheses of homologous ω-aminated 1-methoxyalkyl β-D-glucopyranosides as potential, 53
$N^1$-(*p*-Glycosyloxycinnamoyl)spermidines, synthesis of, 171

Heptofuranose with sodium methoxide: an aldehyde-group participation, the reaction of 2,3:6,7-di-*O*-isopropylidene-5-*O*-methanesulphonyl-L-*glycero*-D-*manno*-, 234

Intramolecular hydrogen-bonding and solvation contributions to the relative stability of the β-furanose form of D-fructose in dimethyl sulfoxide, 159
Isomerization of monosaccharides by alkali, contribution of the reaction pathways involved in the, 13

"Lacto-*N*-biose II" unit, synthesis of three trisaccharides containing the, 201
Lactose derivatives, glycosylation at OH-3′ and OH-4′ of, 105
Lipid intermediate (of glycoprotein biosynthesis), tetrasaccharide-, synthesis, 213

Mannitol and 2-*C*-(hydroxymethyl)-D-mannitol, branched-chain alditols: a convenient synthesis of 6-deoxy-2-*C*-methyl-D-, 252
Monosaccharides by alkali, contribution of the reaction pathways involved in the isomerization of, 13

N.m.r. spectra of methyl 2-deoxy-2-fluoro- and 3-deoxy-3-fluoro-α- and β-D-glucopyranosides, synthesis and, 23
*C*-Nucleoside, 1,3-dimethyl 8-β-D-ribofuranosyl xanthine monohydrate, the crystal structure of the, 35
Nucleotides, f.t.-i.r. spectra of oligo- and poly-, 83

Octyl α-D-glucopyranoside monohydrate and hemihydrate, crystal structures, 1
Oligosaccharides by medium-pressure anion-exchange chromatography on Mono Q, separation of sialyl-, 43

Preparation of 4,6-*O*-benzylidene-2-*C*- and -3-*C*-*p*-tolylsulfonylhex-2-enopyranoside derivatives and a 2-*C*-*p*-tolylsulfonylhex-2-enitol derivative, 189

Reaction of 2,3:6,7-di-*O*-isopropylidene-5-*O*-methanesulphonyl-L-*glycero*-D-*manno*-heptofuranose with sodium methoxide: an aldehyde-group participation, 234
Reaction pathways involved in the isomerization of monosaccharides by alkali, contribution of the, 13
Route to a functionalized anhydro derivative of 1,6-anhydro-β-D-glucopyranose (levoglucosan), a ready, 247

Selective deuteration over Raney nickel in deuterium oxide: 1,6-anhydrohexoses, 151
Separation of sialyl-oligosaccharides by medium-pressure anion-exchange chromatography on Mono Q, 43
Sialyl-oligosaccharides by medium-pressure anion-exchange chromatography on Mono Q, separation of, 43
Spectra of oligo- and poly-nucleotides, F.t.-i.r., 83
Spermidines, synthesis of $N^1$-(*p*-glycosyloxycinnamoyl), 171
Stability of the β-furanose form of D-fructose in dimethyl sulfoxide, intramolecular hydrogen-bonding and solvation contributions to the relative, 159
Syntheses of homologous ω-aminated 1-methoxyalkyl β-D-glucopyranosides as potential β-D-glucosidase inhibitors, 53
Synthesis of antitumor-active 7-*O*-(2,6-dideoxy-2-fluoro-α-L-talopyranosyl)-daunomycinone and -adriamycinone, 69
Synthesis of 5-*O*-(4-deoxy-4-fluoro-β-D-mycaminosyl)tylonolide, 241
Synthesis of 6-deoxy-2-*C*-methyl-D-mannitol and 2-*C*-(hydroxymethyl)-D-mannitol, branched-chain alditols: a convenient, 252
Synthesis of $N^1$-(*p*-glycosyloxycinnamoyl)spermidines, 171
Synthesis of n.m.r. spectra of methyl 2-deoxy-2-fluoro- and 3-deoxy-3-fluoro-α- and β-D-glucopyranosides, 23

Tetrasaccharide-lipid intermediate (of glycoprotein biosynthesis), synthesis, 213
Total synthesis of cyclomaltooctaose and an isomer of cyclomaltohexaose, cyclo-{→6)-[α-D-Glc*p*-(1→4)]$_5$-α-D-Glc*p*-(1→}, 127
-3-*C*-*p*-Tolylsulfonylhex-2-enopyranoside derivatives and a 2-*C*-*p*-tolylsulfonylhex-2-enitol derivative, preparation of 4,6-*O*-benzylidene-2-*C*- and, 189
Tylonolide, synthesis of 5-*O*-(4-deoxy-4-fluoro-β-D-mycaminosyl), 241